Keeping Your Bones Healthy after Childhood Cancer

During childhood and into young adulthood, bone formation usually occurs faster than bone loss, causing bones to grow and become heavier (more dense). As a person gets older, the process of bone removal gradually overtakes bone formation, and bones slowly lose strength as part of the normal aging process. However, loss of bone strength may occur at earlier ages in childhood cancer survivors because of certain cancer treatments. Loss of bone strength may result in a condition known as osteoporosis, which is sometimes referred to as "low bone mineral density."

Osteoporosis: A Silent Disease

Osteoporosis is a disorder resulting from too little new bone formation or too much bone loss, causing bones to become weak. Most people do not have symptoms, especially in the early stages. However, as bones become weaker, the risk for fractures increases. Osteoporosis may occur in any bone, but most commonly affects the wrists, hips, spine, and leg bones.

How is osteoporosis diagnosed?

Although osteoporosis may be suspected based on a patient's symptoms and risk factors, the diagnosis is made by measuring bone density with special x-ray techniques, called DEXA or bone density scans. These scans do not expose patients to large amounts of radiation, and generally take less than 20 minutes to perform.

People who have osteoporosis should discuss treatment options with their healthcare provider. Medications, such as bisphosphonates and calcitonin, are available specifically for the treatment of low bone density. In addition, if you have low levels of male or female hormones, or low levels of growth hormone, you may also benefit from hormone replacement therapy.

What are the risk factors for osteoporosis?

Osteoporosis is more common in people with the following characteristics:

- Female (especially after menopause)
- Family history of osteoporosis
- Caucasian or Asian race
- Small, thin frame
- Older age

The following factors may also increase the risk of osteoporosis:

- Smoking
- Diet low in calcium



Copyright 2008 © Children's Oncology Group All rights reserved worldwide

- Lack of weight-bearing exercise
- Too much caffeine, alcohol, or soda
- A diet high in salt

Additional causes of osteoporosis in people who have had cancer may include:

A history of treatment with:

- Corticosteroids (such as prednisone and dexamethasone)
- Methotrexate
- Radiation to weight-bearing bones (legs, hips, spine)

Conditions resulting from cancer treatment, including:

- Low levels of female or male hormones
- Growth hormone deficiency
- High levels of thyroid hormone
- Chronic graft-versus-host disease requiring prolonged therapy with corticosteroids
- Prolonged periods of inactivity (bed rest)

Other medical treatments, including:

- Certain anticonvulsants (phenytoin and barbiturates)
- Aluminum-containing antacids (such as Maalox[®] or Amphogel[®])
- Medications such as Lupron (used for treatment of early puberty and endometriosis)
- High doses of heparin (used to prevent blood clots), especially with prolonged use
- Cholestyramine (used to control blood cholesterol)

Many of the medications on this list are essential treatments for certain medical conditions. If you are taking any of these medications, do not change your dosage or stop taking your medication without consulting with your healthcare provider.

What lowers the risk of osteoporosis?

Fortunately, there are many things you can do to reduce the risk of osteoporosis. Regular weight-bearing exercise (such as brisk walking, dancing, jazzercise and jogging) helps to develop and maintain healthy bones. Bicycling and swimming are excellent exercises for general fitness, but these are NOT weight-bearing exercises, and they do not help to build strong bones. **Exercises that are especially good for bone health include higher-impact weight-bearing activities, such as hopping, jogging and jumping rope. Resistance exercises, such as light weight lifting, also help to build strong bones** and are especially important for bones of the upper



body, including the arms and shoulders. If you have problems with your heart, or have painful bones or joints, **be sure to discuss your individual health status and cancer treatment history with your healthcare provider before starting any new exercise program.**

A diet high in calcium also is important in preventing osteoporosis. Most healthcare professionals recommend 1000-1500 mg a day, which means a diet rich in dairy products (milk, cheese, yogurt) and leafy green vegetables. Talking with a dietitian may help you design a healthy diet. Over- the-counter calcium supplements also may be useful. See Tables 1 and 2 for recommendations for calcium intake. Additional information about calcium-rich diets is available at www.nationaldairycouncil.org.

Vitamin D is needed in order to absorb calcium. Skin makes this vitamin naturally when exposed to sunlight. Many dairy products also contain vitamin D. In general, 200 units of Vitamin D is recommended daily. You should not take more than 800 units of Vitamin D per day. Taking too much vitamin D may be harmful, so it's best to check with your healthcare provider before taking any vitamin D supplements.

What screening is recommended?

After reviewing your treatment history and risk factors, your healthcare provider can advise you regarding the need for bone density testing. For those at risk, a baseline bone density scan is recommended for childhood cancer survivors when they enter long-term follow-up (2 or more years after completion of therapy). Follow-up scans may be needed for ongoing monitoring of bone density in some patients.

Table 1: Recommendations for Adequate Dietary Calcium Intake in the United States

Age	Recommended Calcium Intake
1 - 3 years	500 mg per day
4 - 8 years	800 mg per day
9 - 18 years	1300 mg per day
19 - 50 years	1000 mg per day
50 - 70+ years	1200 mg per day

(from the Food and Nutritional Board, Institute of Medicine, 1997)



Table 2: Common Foods that are Good Sources of Calcium

Food	Serving Size	Calcium Content	Number of servings to equal calcium in 1 cup low-fat milk
Dairy foods			
Whole milk	1 cup (244 g)	246 mg	1.0
Low-fat (1%) milk	1 cup (244 g)	264 mg	1.0
Nonfat milk	1 cup (245 g)	223 mg	1.2
Yogurt, nonfat, fruit variety	6 oz (170 g)	258 mg	1.0
Frozen yogurt, vanilla, soft serve	1/2 cup (72 g)	103 mg	2.6
Cheese	1 1-oz slice (28 g)	202 mg	1.3
Cheese, pasteurized, processed	1 3/4-oz slice (21 g)	144 mg	1.8
Cheese, ricotta, part skim milk	1/2 cup (124 g)	337 mg	0.7
Nondairy foods			
Salmon, sockeye canned, drained, with bones	3 oz (85 g)	203 mg	1.3
Tofu, firm, prepared with calcium sulfate and magnesium chloride	1/2 cup (126 g)	204 mg	1.3
White beans, cooked, boiled	1 cup (179 g)	161 mg	1.6
Broccoli, cooked	1 cup, chopped (156 g)	62 mg	4.3
Collards, cooked, boiled, drained	1 cup, chopped (190 g)	266 mg	1.0
Baked beans, canned	1 cup (253 g)	127 mg	2.1
Tomatoes, canned, stewed	1 cup (255 g)	87 mg	3.0
Foods fortified with calcium		1	
Calcium-fortified orange juice	1 cup (240 ml)	300 mg	0.9
Selected fortified breakfast cereals	3/4 - 1 cup (30 g)	100 mg	2.6
Instant oatmeal, fortified, plain, prepared with water	1/2 cup (117 g)	65 mg	4.1
English muffin, plain, enriched, with calcium propionate	1 muffin (57 g)	99 mg	2.7
Calcium-fortified soy milk	1 cup (240 ml)	200 – 500 mg	0.5 – 1.3

(from the U.S. Department of Agriculture Research Service, 2005)



Written by Julie Blatt MD, Division of Pediatric Hematology-Oncology, University of North Carolina, Chapel Hill, NC, and Lillian R. Meacham, MD, Division of Pediatric Endocrinology, Emory University, Children's Healthcare of Atlanta, Atlanta, GA

Reviewed by Charles Sklar MD, Melissa M. Hudson MD, Debra L. Friedman MD, Joan Darling PhD, Wendy Landier CPNP, and Sarah Bottomley MN, RN, CPNP, CPON

Additional health information for childhood cancer survivors is available at www.survivorshipguidelines.org

Note: Throughout this *Health Links* series, the term "childhood cancer" is used to designate pediatric cancers that may occur during childhood, adolescence, or young adulthood. *Health Links* are designed to provide health information for survivors of pediatric cancer, regardless of whether the cancer occurred during childhood, adolescence, or young adulthood.

Disclaimer and Notice of Proprietary Rights

Introduction to Late Effects Guidelines and Health Links: The "Long-Term Follow-Up Guidelines for Survivors" of Childhood, Adolescent, and Young Adult Cancers" and accompanying "Health Links" were developed by the Children's Oncology Group as a collaborative effort of the Late Effects Committee and Nursing Discipline.

For Informational Purposes Only: The information and contents of each document or series of documents made available from by the Children's Oncology Group relating to late effects of cancer treatment and care or containing the title "Long-Term Follow-Up Guidelines for Survivors of Childhood, Adolescent, and Young Adult Cancers" or the title "Health Link", whether available in print or electronic format (including any digital format, e-mail transmission, or download from the website), shall be known hereinafter as "Informational Content". All Informational Content is for informational purposes only. The Informational Content is not intended to substitute for medical advice, medical care, diagnosis or treatment of the aphysician or health care provider.

To cancer patients (if children, their parents or legal guardians): Please seek the advice of a physician or other qualified health provider with any questions you may have regarding a medical condition and do not rely on the informational Content. The Children's Oncology Group is a research organization and does not provide individualized medical care or treatment.

To physicians and other healthcare providers: The Informational Content is not intended to replace your independent clinical judgment, medical advice, or to exclude other legitimate criteria for screening, health counseling, or intervention for specific complications of childhood cancer treatment. Neither is the Informational Content intended to exclude other reasonable alternative follow-up procedures. The Informational Content is provided as a courtesy, but not intended as a sole source of guidance in the evaluation of childhood cancer survivors. The Children's Oncology Group recognizes that specific patient care decisions are the prerogative of the patient, family, and healthcare provider.

No endorsement of any specific tests, products, or procedures is made by Informational Content, the Children's Oncology Group, or affiliated party or member of the Children's Oncology Group.

No Claim to Accuracy or Completeness: While the Children's Oncology Group has made every attempt to assure that the Informational Content is accurate and complete as of the date of publication, no warranty or representation, express or implied, is made as to the accuracy, reliability, completeness, relevance, or timeliness of such Informational Content. No Liability on Part of Children's Oncology Group and Related Parties' Agreement to Indemnify and Hold Harmless the Children's Oncology Group and Related Parties' No liability is assumed by the Children's Oncology Group or any affiliated party or member thereof for damage resulting from the use, review, or access of the Informational Content. You agree to the following terms of indemnification: (i) "Indemnified Parties" include authors and contributors to the Informational Content, all officers, directors, representatives, employees, agents, and members of the Children's Oncology Group and affiliated organizations; (ii) by using, reviewing, or accessing the Informational Content, you agree, at your own expense, to indemnify, defend and hold harmless Indemnified Parties from any and all losses, liabilities, or damages (including attorneys' fees and costs) resulting from any and all claims, causes of action, suits, proceedings, or demands related to or arising out of use, review or access of the Informational Content.

Proprietary Rights: The Informational Content is subject to protection under the copyright law and other intellectual property law in the United States and worldwide. The Children's Oncology Group retains excursive copyright and other right, title, and interest to the Informational Content and claims all intellectual property rights available under law. You hereby agree to help the Children's Oncology Group secure all copyright and intellectual property rights for the benefit of the Children's Oncology Group by taking additional action at a later time, action which could include signing consents and legal documents and limiting dissemination or reproduction of Informational Content.



Copyright 2008 © Children's Oncology Group All rights reserved worldwide